

# **Anthropometric Measurement of the developing Human Fetus**

## **Introduction:**

The scientific study of the measurements and proportions of the human body is known as Anthropometry. These studies are generally used by clinicians and pathologists for adequate assessments of the growth and development of the fetus at any specific point of gestational maturity. Fetal height, Fetal Weight and head circumference (HC), Crown to Rump length(CR), dermatological observations like skin thickness etc are measured individually to assess the growth and development of the organs and the fetus as a whole. It is also an essential and important parameter for assessing the fetus to adapt to its new environment. (1)

Another important factor that contributes towards the anthropometric measurement of the human fetal growth is the maternal nutrition and maternal well-being. Malnutrition as already established by WHO is a global serious health problem not only in adults but in pregnant and lactating mothers too. In Africa and South Asia 27-50% of women of reproductive age are underweight resulting in 30 million low birth weight babies. (2)

For decades, the topic of question pertaining to crown rump length (CR), crown-heel length (CH), head circumference (HC) with respect to the body weight of human fetus at different time periods of gestation has baffled many developmental researchers and biostatisticians.)These variations of biological human parameters are all based on linear curves based on human fetuses between 9 to 28 weeks of gestation. (3,4,5,6,7, 8)

## **Co-relation of fetal weight and fetal growth**

Body weight for example is an important function and parameter for growth with respect to gestational age of the fetus. There will be great variations in the body weight of a 16 weeks old fetus. The weight will not be constant for every fetus and will vary from individual to individual. The weight will not be constant in other words. Therefore, rather than an appropriate or standard value, a range can be specified like 90 to 100 grams. This amount of variations applies to all other anthropometric measurements. Often, the scientific world cover up their ignorance by stating that the rate of growth of particular human fetus depends upon its intrinsic growth potential environment provided by the normal mother and is a visible function of the genetic potential.(9)

The fetal growth is not an individual growth and is dependent on the composite growth of the organs which are constitutive of the organs. (10, 11, 12, 13) Growth of the individual organs is controlled by the genetic potential, the environment provided by the mother and by the fetus itself. Scientists have or are trying to determine such relationships by a number of works. (14, 15, 16)

Streeter, Schults et al., all studied the fetal dimensions obtained from spontaneous abortions and pathological pregnancies on mainly formed and fixed specimens. (17) The growth of an organ from inception to a definitive functional stage is dependent on the integrated function of the whole organism which depends on a number of parameters such as nucleic acid content of the cells in the organ being one of the important factors. Functioning of an organ is important for the development of the whole organ and is therefore related to the whole organism. (18)

A Nigerian study showed that the birth weight of the human fetus also depends upon the size and weight of the mother including her height and weight. (19) Further a Polish study reported a similar reported that some measurements like the ear height, muscular strength of the shoulders, skin fold thickness, mandibular breadth including the height of the upper and the lower limbs can be co-related to the mother also. (20) Similar observations were also reported by Gueri et al. (21)

One of the first original and unique works to be conducted on the anthropometric measurement of the human fetus in the Indian context was conducted by a group of scientists in Calcutta between 1977 to 1987 under the supervision of K.L.Mukherjee, a stalwart in the field of medical biochemistry in Institute of Post Graduate Medical Education and Research. (22)

The researchers divided the fetuses into 6 groups of A, B, C, D, E and F with a difference of 4 weeks of gestation period among the 6 groups. Group A had 90 fetuses of 9-12 weeks of gestation and the weight varied between 1-14 grams. Group B had 337 fetuses, age 13-16 weeks of gestation with weight variation between 15-105 grams whereas the third group had 435 fetuses of 17-20 weeks of gestation with a weight range between 106-310 grams. Group D consisted of 531 fetuses of 21-24 weeks and weight between 331-640 grams and Group E had fetuses of the age range 25-28 weeks of gestation and weight 640-1070grams. The last two groups F and G had fetuses with gestation period between 29-32 and 33-36 weeks. All aborted fetuses were collected after permission from the Institute Ethics committee followed with donor consent form with primary objective being the aborted mother's health and safety. (22)

#### **Liver growth:**

Researchers observed that the liver weight is directly proportional to the body weight. At 8-12 weeks of gestation, liver is a bigger organ which forms 4-5-5.5% of the total body weight and protrudes through the abdominal wall. From 13 to 32 weeks of gestation, it forms 3.4% to 4.0 % of the total body weight. (23)The liver weight hence forms a more or less constant proportion of the total body weight of the fetus. (24)

#### **Growth of the lung:**

Although in adult life, the lung is the only major respiratory organ, in case of fetal life such is not the case though the fetal lung is known to expand and contract in the last phase of development. Both the weight of the right and left lungs are normally assessed at different periods of gestation and is expressed as a function of the total body weight. (22)

An irregular graph was observed by K.L.Mukherjee and his group instead of the standard normal linear graph after plotting a graph of weight of the lungs expressed as gm/kg of body weight against the body weight. This relationship was observed from fetuses weighing 350 grams to 850 grams after which the rate of the growth became uniformly proportional again. (22)

#### **Brain and Central nervous System:**

Brain and the central nervous system are the two most important components of the fetus. Further analysis by this same group involved the CNS upto the medulla at the level of the 2<sup>nd</sup> cervical vertebra. (22) The process of analyzing the fetal brain and the CNS involved dissecting out the whole mass of brain tissues followed by decantation of the CNS of a 15g gram of fetus with 8.5 weeks of gestational period. The brain at this time had already assumed the appearance of primary divisions and flexures, and the prosen, messen and rhombocephalon already gave rise to the different brain derived constituents like rhinocephalon, corpora striata, cerebral cortex, hypo and epithalamus and pons medulla to a less differentiation extent. (22) The growth of the fetal brain from this time onwards was proportional to the body weight although some brains at this stage can weigh between 20% to even 12% or 13% of the body weight by and large. Scientists are still yet to find an explanation for this. (22)

#### **Kidney and the Adrenal Glands:**

In early gestational period, the weight of the adrenal glands outweighs even the metanephric kidneys and is comparatively a larger organ. After 10<sup>th</sup> week of gestation, the kidney grows at a much

rapid rate than the adrenal glands. Hence with increase in gestational time, i.e. by 12<sup>th</sup> week of gestation both the kidneys and adrenal glands measure the same. However, post 12 weeks the kidneys measure more than the adrenal glands. However, the Adrenal gland is a larger organ in the fetus than the adult. The same group of researchers further observed that with the increase of fetal age, the adrenal glands also weigh more as observed in 90 human fetuses. However, the rate of increase is not uniform and varies throughout the fetal growth. (25, 26)

#### **Human fetal testes:**

The growth of the fetal testes is not uniform as revealed through various other studies. Further, it was found that the weight of the right testes was more than the weight of the left testes which was according to anatomical laws. Exceptions were however noticed in some of the cases as reported by K.L.Mukherjee and his group. Normally like all other organs the growth of the testes including its weight also increases with increase in the gestational period. The research group further through their graph plot, examined that the growth of testes was not uniform with proportional growth initially which soon flattened to increase with different slopes and spikes consistently throughout the whole length of the gestational period. Further the weight of the human testes marked as mg/100 gram of body weight was investigated. It was observed that there was a steep decline in the early gestation period from about 200 mg/100 gram of body weight to 60 mg/ 100 gram of body weight when the fetal weight was about 1.5 grams to 20 grams. In case of a 1.6 kilogram fetus, the testes weighed only 20 mg/100 gram of body weight. This decline was however not maintained uniformly. (22)

#### **Growth of the Human fetal ovaries:**

A steep decline in the ovarian weight in the early gestational period was observed though it was not a uniformly maintained decline. With increase in the gestational time, progressive weight of the ovaries were found and in most cases the weight of the ovaries were identical to the weight of the fetuses although some exceptions were observed by the group. (22)

#### **Fetal Thymus growth:**

At 8 weeks of gestation when the fetus weighed 1 gram, the thymus could not be detected. In many of the 39 fetuses weighing around 1.3-14.7 grams, the thymus tissues could not be dissected by the group especially in the smaller fetuses. Fetuses weighing more than or equal to 5 grams could be detected. Plotting a graph it was observed that thymus organ formed 52 mg/100 gram of body weight in case of a 5 gram fetus. Further study on 28 fetuses weighing 15 to 100 grams revealed the thymic weight to be 77 mg per 100 grams of the body weight. The relative growth of the thymus was more in this group compared to all the earlier observations. A further group including 39 fetuses weighing between 100-300 grams showed a fetal thymic weight between 136 mg/100 gram to 77 mg/ 100 gram. (22) In fetuses up to 28 weeks, it was observed by scientists that the fetal thymic weight was the highest and was in contrast to many other organs like brain, liver which constitute more or less constant proportion of the body weight with very few exceptions. Therefore, it was inferred that with increase in the gestational period, the thymic weight also increase although exceptions were observed. (27)

#### **Conclusion:**

Growth and development throughout the fetal life are two most important factors which determine the growth rate of each individual and their specific organs. This process of maturation and development of the organs are observed in post natal life also. With increase in gestational time, the fetal organs also grow in progression to the body weight, the phenomenon which is still not

understood clearly by many researchers. Some believe that genetic potentiality of the different endocrine organs related to the growth and various other unidentified processes mediate the

#### **References:**

1. World Health Organization. Public health aspect of low birth weight, third report of the expert committee on maternal and child health (WHO), Geneva. 1961.
2. Monika Blössner Mercedes de Onis, Environmental Burden of Disease Series, No. 12 Malnutrition Quantifying the health impact at national and local levels., World Health Organization Nutrition for Health and Development Protection of the Human Environment Geneva 2005.
3. Lubchenco LO, Hansman C, Dressler M, Boyd E. Intrauterine growth as estimated from liveborn birth-weight data at 24 to 38 weeks of gestation. *Pediatrics*. 1963;32:793–800.
4. Lubchenco LO, Hansman C, Boyd E. Intrauterine growth in length and head circumference as estimated from live births at gestational ages from 26–42 weeks. *Pediatrics*. 1966;37:403–8.
5. Usher R, McLean F. Intrauterine growth of live-born Caucasian infants at sea level; standards obtained from measurements in 7 dimensions of infants born between 25 and 44 weeks of gestation. *J Pediatr*. 1969;74:901–10.
6. Berger GS, Edelman DA, Kerenyi TD. Fetal crown-rump length and biparietal diameter in the second trimester of pregnancy. *Am J Obstet Gynecol*. 1975;122:9–12.
7. Birkbeck JA, Billewicz WZ, Thomson AM. Human fetal measurements between 50 and 150 days of gestation in relation to crown-heel length. *Ann Hum Biol*. 1975;2:173–8.
8. Drumm JE, Clinch J, Mackenzie G. The ultrasonic measurement of fetal crown-rump length as a method assessing gestational age. *Br J Obstet Gynaecol*. 1976;83:417–21.
9. Bhatia BD, Tyagi NK. Birth weight; relationship with other fetal anthropometric parameters. *Indian Pediatr*. 1984;21:833–8.
10. Campbell S. The prediction of fetal maturity by ultrasonic measurement of the biparietal diameter. *J Obstet Gynaecol Br Commonw*. 1969;76:603–9.
11. Parekh UC, Pherwani A, Udani PM, Mukherjee S. Brain weight and head circumference in a fetus, infant and children of different nutritional and socio-economic groups. *Indian Pediatr*. 1970;7:347–58.

12. Prema LN, Nagaswamy S, Raju VB. Fetal growth as assessed by anthropometric measurements. *Indian Pediatr.* 1974;11:803–10.
13. Brenner WE, Edelman DA, Hendricks CH. A standard of fetal growth for the United States of America. *Am J Obstet Gynecol.* 1976;126:555–64.
14. Hern WM. Correlation of fetal age and measurements between 10 and 26 weeks of gestation. *Obstet Gynecol.* 1984;63:26.
15. Mukherjee B, Mitra SC, Gunasegaran JP. Fetal crown-rump length and body weight at different gestational periods. *Indian J Med Res.* 1986;83:495–500.
16. Sood M, Hingorani V, Kashyap N, Kumar S, Berry M, Bhargava S. Ultrasonic measurement of fetal parameters in normal pregnancy and in intrauterine growth relation. *Indian J Med Res.* 1988;87:453.
17. Schultz AH. Fetal growth in man. *Am J Phys Anthropol.* 1923;6:389–99.
18. Schultz AH. Fetal growth in man and other primates.
19. Baird SD. The epidemiology of prematurity. *J Paedit* 1965;65: 909-923.
20. Wolanski N, Spruch HC. Maternal regulation of intrauterine life: Anthropology of maternity. Charles University, Prague. 1977; 109-117.
21. Gueri M, Justin P, Sorhaindo B. Anthropometric assessment of nutritional status in pregnant women : A reference table of weight for height by a week of pregnancy. *A J Clin Nutr* 1982; 35:609-616.
22. Bhattacharya, Niranjana, Stubblefield, Phillip G: Editors, Human Fetal Growth and Development, First and Second Trimesters, Chapter No: 6, Anthropometric Measurement of the Human Fetus, Page Nos: 67-83, Springer International Publishing, Switzerland, ISBN: 978-3-319-14.
23. Bhattacharya, Niranjana, Stubblefield, Phillip G: Editors, Human Fetal Growth and Development, First and Second Trimesters, Chapter No: 26, Growth and Development of Human Liver and Its Future Implications, Gopal Krishna Dhali, Gurubasava Lakamaji, Page Nos: 320, Springer International Publishing, Switzerland, ISBN: 978-3-319-14.
24. Bhattacharya, Niranjana, Stubblefield, Phillip G: Editors, Human Fetal Growth and Development, First and Second Trimesters, Chapter No: 25, Embryonic Development of Human Liver and Its Future Implications, Abhijeet Chaudhuri, Page Nos: 333, Springer International Publishing, Switzerland, ISBN: 978-3-319-14.

25. Bhattacharya, Niranjana, Stubblefield, Phillip G: Editors, Human Fetal Growth and Development, First and Second Trimesters, Chapter No: 10, Mechanism of Rejection of Human Fetal Adrenal Cortex, Chameli Ganguly Bimal Samanta, Gitanjali Guha Thakurata, Chaitali Bhattacharya, Reba Bhattacharya, K. L. Mukherjee, Niranjana Bhattacharya, Page Nos: 135-159, Springer International Publishing, Switzerland, ISBN: 978-3-319-14.

26. Bhattacharya, Niranjana, Stubblefield, Phillip G: Editors, Human Fetal Growth and Development, First and Second Trimesters, Chapter No: 31, Development of renal tissues in first twenty weeks, Rajendra Pandey, Page Nos: 319-328, Springer International Publishing, Switzerland, ISBN: 978-3-391-395.

27. Bodey B, Kaiser HE. Development of Hassall's bodies of the thymus in humans and other vertebrates (especially mammals) under physiological and pathological conditions: immunocytochemical, electron microscopic and in vitro observations. *In Vivo*. 1997; 11:61-85.

**See Also:**

Anthropometry

Prenatal Development

**External Links:**

Anthropometric measurements: [http://www.thl.fi/publications/ehrm/product2/part\\_iii5.htm](http://www.thl.fi/publications/ehrm/product2/part_iii5.htm)

Anthropometric Indicators Measurement Guide:

<http://www.fantaproject.org/sites/default/files/resources/anthropometry-2003-ENG.pdf>

**Further Readings:**

[http://www.micronutrient.org/nutritiontoolkit/ModuleFolders/3.Indicators%5CAnthropometry%5CResources%5CPhysical\\_status\\_-\\_The\\_use\\_and\\_interpretation\\_of\\_anthropometry%5CChapter\\_3\\_\(Preg\\_women\).pdf](http://www.micronutrient.org/nutritiontoolkit/ModuleFolders/3.Indicators%5CAnthropometry%5CResources%5CPhysical_status_-_The_use_and_interpretation_of_anthropometry%5CChapter_3_(Preg_women).pdf)

<https://www.cdc.gov/nchs/data/nhanes/nhanes3/cdrom/nchs/manuals/anthro.pdf>